

Searching for dark matter and exotic physics with atomic clocks in space and on the ground

Benjamin M. Roberts

(University of Queensland, Australia)

G. Blewitt, C. Dailey, M. Pospelov, G. Panelli, and A. Derevianko

(UNR, Reno; Perimeter Institute)

P. Delva, A. Hees, E. Savalle, P. Wolf

(SYRTE, Observatoire de Paris)

& the Optical clock, link, comb, & cavity teams of NPL, PTB, & SYRTE


Scientific and Fundamental Aspects of GNSS
7th International Colloquium, European Space Agency (ESA)
ETH Zürich, Switzerland




4–6 September 2019

Dark Matter: What is it?

- $\sim 25\%$ of Universe energy budget
(cf $\sim 5\%$ for “normal” matter)
- Possible mass range: ~ 90 orders-of-magnitude:



[image: wmap.gsfc.nasa.gov]



[• US Cosmic Visions report, arXiv:1707.04591]

(context: $m_{\text{Earth}} \sim 10^{60} \text{ eV}$ $m_{\text{electron}} \sim 10^6 \text{ eV}$)


⇒ Wide range of possibilities: requires large range of searches

Dark Matter Clumps: (Topological Defects)

- Ultralight ($m_\phi \ll eV$) \Rightarrow high occupation number
Many possibilities: Here: TDs

Topological Defects

- monopoles, strings, walls,
- Defect width: $d \sim 1/m_\phi$
- Earth-scale object: $m_\phi \sim 10^{-14} \text{ eV}$




$$\text{Inside: } \phi^2 \rightarrow A^2, \quad \text{Outside: } \phi^2 \rightarrow 0$$

Dark matter: Gas of defects

- DM: galactic speeds: $v_g \sim 10^{-3}c$
- Collisions offer chance for lab detection


- Vilenkin '85, Coleman '85, Lee '89, Kibble '80, ...
- Derevianko, Pospelov, Nature Phys. 10, 933 (2014).



Transients: 3D(+) parameter space

Two time-scales

- \mathcal{T} – time between events
 - Given by number density
- $\tau_{\text{int}} = v/d$ – interaction duration
 - For TDs: $d \sim 1/m_\phi$ (free in general)
- For transients: must have $\tau_{\text{int}} \ll \mathcal{T}$



$$\phi_0^2 = [\rho_{\text{DM}}] v_g d \mathcal{T},$$

$\rho_{\text{DM}} \simeq 0.4 \text{ GeV/cm}^3$: local DM density

And (hopefully) some non-gravitational coupling

- Various possibilities. Here: (quadratic) scalar:
- $\mathcal{L}_{\text{int}} \sim \phi^2 (a \bar{\psi} \psi + b F_{\mu\nu}^2 + \dots)$

See, e.g.: Olive, Pospelov, Phys. Rev. D **65**, 085044 (2002); Derevianko, Pospelov, Nature Phys. 10, **933** (2014); Pustelny, Kimball, Pankow, Ledbetter, Wlodarczyk, Wcislo, Pospelov, Smith, Read, Gawlik, Budker, Ann.Phys. **525**, 659 (2013)
Stadnik, Flambaum, PRL **114**, 161301 (2013).

Hees, Guena, Abgrall, Bize, Wolf, Phys. Rev. Lett. **117**, 061301 (2016).

Arvanitaki, Graham, Hogan, Rajendran, Van Tilburg (2016); Stadnik, Flambaum (2016), ...

Variation of fundamental constants

- Here: (quadratic) scalar: $\mathcal{L}_{\text{int}} \sim \phi^2(a\bar{\psi}\psi + bF_{\mu\nu}^2 + \dots)$

\implies transient additions to *effective values* of fundamental constants

$$\alpha^{\text{eff}}(r, t) = \alpha \left(1 + \frac{\phi^2(r, t)}{\Lambda_\alpha^2} \right), \quad m_f^{\text{eff}}(r, t) = m_f \left(1 + \frac{\phi^2(r, t)}{\Lambda_f^2} \right),$$

\implies shifts in energy levels \implies shifts in clock frequencies

$$\frac{\delta\omega(r, t)}{\omega_0} = K_\alpha \frac{\delta\alpha(r, t)}{\alpha} = \phi^2(r, t) \frac{K_\alpha}{\Lambda_\alpha^2}$$


Monitor Atomic Clocks

- Clocks: lock frequency to atomic transition
- Shift $\delta\omega$ occurs only when ϕ non-zero (inside DM object)
- Monitor atomic frequencies using atomic clocks
- Parameterised in with Λ “energy scale” (\sim inverse coupling strength)

Shift in atomic clock frequencies


Monitor Atomic Clocks

- Temporary frequency shift \rightarrow bias (phase) build-up
- Initially synchronised clocks become desynchronised



Signal v. noise?

- Transient signal:
looks essentially like any outlier
- i.e. what is the specific DM signature?




- Derevianko, Pospelov, Nat. Phys. 10, 933 (2014).

Global network of precision devices

Network of separated atomic clocks


- DM expected to move at \sim galactic speeds
- Correlated signal propagation through network, $v \sim 300 \text{ km/s}$
- \vec{v} encoded in time-delay, ordering: $\Delta t \sim \text{seconds} - \text{minutes}$
- Also: multiple clock-types in network, each has different K_α (relative sensitivities prediction of theory)



- Clocks: Derevianko, Pospelov, Nature Phys. 10, 933 (2014).
- Magnetometer: Pospelov, Pustelny, Ledbetter, Kimball, Gawlik, Budker, Phys.Rev.Lett. 110, 021803 (13).



GPS: 50,000 km DM observatory

- 32 satellite clocks (Rb/Cs), ~ 16 years of high-quality data
- Also several H-maser ground-based clocks.
- Data from JPL: (sideshow.jpl.nasa.gov/pub/jpligsac/)
 - 30s sampled data; 0.01–0.1 ns precision
- Correlated, directional signal, with $v_g \sim 300$ km/s




DM Walls: Initial GPS search/limits


- Thin wall: brief (< 30 s) frequency excursion



- \vec{v} encoded in time-delay and signal ordering: $\Delta t \sim$ minutes




Apply $S_{\text{cut}} \implies$



Results: Limits - Λ_α (photon)

- Exclusion plot (assume coupling to photon dominates)



- BMR, Blewitt, Dailey, Murphy, Pospelov, Rollings, Sherman, Williams, Derevianko, Nature Comm. **8**, 1195 (2017).



Sr: Wcislo, Morzynski, Bober, Cygan, Lisak, Ciurylo, Zawada, Nat. Astron. **1**, 9 (2016).

Astro: Olive, Pospelov, Phys. Rev. D. **77**, 43524 (2008).

Results: Limits - fermion masses

Combine Rb, Cs, and Sr (optical)

- Three different combo's of three couplings



- BMR, Blewitt, Dailey, Murphy, Pospelov, Rollings, Sherman, Williams, Derevianko, Nature Comm. **8**, 1195 (2017).

Sr: Wcislo, Morzynski, Bober, Cygan, Lisak, Ciurylo, Zawada, Nat. Astron. **1**, 9 (2016).

Astro: Olive, Pospelov, Phys. Rev. D. **77**, 43524 (2008).

Search for domain wall dark matter with atomic clocks on board global positioning system satellites

Benjamin M. Roberts¹, Geoffrey Blewitt^{1,2}, Conner Dailey¹, Mac Murphy¹, Maxim Pospelov^{3,4},
Alex Rollings¹, Jeff Sherman⁵, Wyatt Williams¹ & Andrei Derevianko¹

Search for transient ultralight dark matter signatures with networks of precision measurement devices using a Bayesian statistics method

B. M. Roberts, G. Blewitt, C. Dailey, and A. Derevianko.

Department of Physics, University of Nevada, Reno 89557, USA


Applying matched-filter technique to the search for dark matter transients with networks of quantum sensors

G. Panelli, B. M. Roberts,* and A. Derevianko



Existing constraints & Discovery frontiers

3D Parameter space:



$$\frac{\delta\omega}{\omega} = K_\alpha \frac{\delta\alpha}{\alpha} = \frac{K_\alpha}{\Lambda_\alpha^2} \phi_0^2$$

- Shown is full \mathcal{T} parameter space (fixed d)

GPS: BMR, Blewitt, Dailey, Murphy, Pospelov, Rollings, Sherman, Williams, Derevianko, *Nature Comm.* **8**, 1195 (2017).

2016: Wcislo, Morzynski, Bober, Cygan, Lisak, Ciurylo, Zawada, *Nat. Astro.* **1**, 00009 (2016).

2018: Wcislo, Ablewski, Beloy, Bilicki, Bober, Brown, Fasano, Ciurylo, Hachisu, Ido, Lodewyck, Ludlow, McGrew, Morzynski, Nicolodi, Schioppo, Sekido, Le Targat, Wolf, Zhang, Zjawin, Zawada, *Sci. Adv.* **4**, 4869 (2018).

Also: *Astro*: Olive, Pospelov, *Phys. Rev. D* **77**, 043524 (2008).

Number density/wait time

- large \mathcal{T} : need long T_{obs}

Sensitivity: Λ_α ($\delta\alpha$)

- Need: small σ_y , large K_X

Size: d (duration τ_{int})

- Existing: only $\lesssim 10^4$ km
- Need: long-term stability

European fibre-linked optical clock network



Fibre network

- High-accuracy long-distance clock-clock (atom-atom) comparisons
- Different clocks: Hg/Sr/Yb⁺
- ~ Days – weeks synchronous running




- Sensitivity: ✓ ($\delta\alpha, \Lambda$)
limited only by clocks:
 $Sr-Sr: \delta\omega/\omega \sim 3 \times 10^{-17}$ at 1000s
 - Long observation time: ✓ (\mathcal{T})
 - Long-term stability: ✓ (d)
-
- Lisdat *et al.* (PTB, LNE-SYRTE), Nature Commun. **7**, 12443 (2016).
 - Delva *et al.* (PTB, SYRTE, NPL, ..), Phys. Rev. Lett. **118**, 221102 (2017).

Transient variation of fine-structure constant

Orders-of-magnitude improvement: especially for large objects (τ)

- $\delta\alpha(\tau)/\alpha \lesssim 5 \times 10^{-17}$ @ $\tau = 10^3$ s, & $\mathcal{T} = 1$ hr
- $\delta\alpha(\tau)/\alpha \lesssim 4 \times 10^{-15}$ @ $\tau = 10^4$ s, & $\mathcal{T} = 45$ hr



• arXiv:1907.02661


2016: Wcisło, Morzynski, Bober, Cygan, Lisak, Ciuryło, Zawada, Nat. Astro. 1, 10009 (2016).

2018: Wcisło, Ablewski, Beloy, Bilicki, Bober, Brown, Fasano, Ciuryło, Hachisu, Ido, Lodewyck, Ludlow, McGrew, Morzynski, Nicolodi, Schioppo, Sekido, Le Targat, Wolf, Zhang, Zjawin, Zawada, Sci. Adv. 4, 4869 (2018).

Topological defect dark matter

Assume DM is made from Topological Defects:

$$\phi_0^2 = \hbar c \rho_{\text{DM}} v_g \mathcal{T} d , \quad \mathcal{T} = \frac{\rho_{\text{inside}}}{\rho_{\text{DM}}} \frac{d}{v_g}$$



- nb: GPS results (orange): go up to $\mathcal{T} \sim 10 \text{ yrs} \sim 10^5 \text{ hrs}$

$$\implies \Lambda_\alpha^2(\mathcal{T}, d) > \frac{\hbar c \alpha \rho_{\text{DM}} v_g \mathcal{T} d}{|\delta \alpha_0(\mathcal{T}, \tau_{\text{int}})|} .$$

GPS: BMR, Blewitt, Dailey, Murphy, Pospelov, Rollings, Sherman, Williams, Derevianko, **Nature Comm.** **8**, 1195 (2017).

2016: Wcislo, Morzynski, Bober, Cygan, Lisak, Ciurylo, Zawada, **Nat. Astro.** **1**, 00009 (2016).

2018: Wcislo, Ablewski, Bely, Bilicki, Bober, Brown, Fasano, Ciurylo, Hachisu, Ido, Lodewyck, Ludlow, McGrew, Morzynski, Nicolodi, Schioppo, Sekido, Le Targat, Wolf, Zhang, Zjawin, Zawada, **Sci. Adv.** **4**, 4869 (2018).

arXiv:1907.02661

Search for transient variations of the fine structure constant and dark matter using fiber-linked optical atomic clocks

B. M. Roberts,^{1,*} P. Delva,¹ A. Al-Masoudi,² A. Amy-Klein,³ C. Bærentsen,¹ C. F. A. Baynham,⁴ E. Benkler,² S. Bilicki,¹ S. Bize,¹ W. Bowden,⁴ J. Calvert,¹ V. Cambier,¹ E. Cantin,^{1,3} E. A. Curtis,⁴ S. Dörscher,² M. Favier,¹ F. Frank,¹ P. Gill,⁴ R. M. Godun,⁴ G. Grosche,² C. Guo,¹ A. Hees,¹ I. R. Hill,⁴ R. Hobson,⁴ N. Huntemann,² J. Kronjäger,⁴ S. Koke,² A. Kuhl,² R. Lange,² T. Legero,² B. Lipphardt,² C. Lisdat,² J. Lodewyck,¹ O. Lopez,³ H. S. Margolis,⁴ H. Álvarez-Martínez,^{1,5} F. Meynadier,^{1,6} F. Ozimek,⁴ E. Peik,² P.-E. Pottie,¹ N. Quintin,⁷ C. Sanner,^{2,†} L. De Sarlo,¹ M. Schioppo,⁴ R. Schwarz,² A. Silva,⁴ U. Sterr,² Chr. Tamm,² R. Le Targat,¹ P. Tuckey,¹ G. Vallet,¹ T. Waterholter,² D. Xu,¹ and P. Wolf^{1,‡}

¹SYRTE, Observatoire de Paris, Université PSL, CNRS,

Sorbonne Université, LNE, 61 avenue de l'Observatoire, 75014 Paris, France

²Physikalisch-Technische Bundesanstalt, Bundesallee 100, 38116 Braunschweig, Germany

³Laboratoire de Physique des Lasers, Université Paris 13, Sorbonne Paris
Cité, CNRS, 99 Avenue Jean-Baptiste Clément, 93430 Villejuif, France

⁴National Physical Laboratory, Hampton Road, Teddington TW11 0LW, United Kingdom

⁵Sección de Hora, Real Instituto y Observatorio de la Armada, San Fernando, Spain

⁶Bureau International des Poids et Mesures, BIPM, Pavillon de Breteuil, 92312 Sèvres, France

⁷Réseau National de télécommunications pour la Technologie,
l'Enseignement et la Recherche, 23–25 Rue Daviel, 75013 Paris, France

(Dated: July 10, 2019)



Systèmes de Référence Temps-Espace



Some new directions

Signal propagation

- DM couples to electromagnetism: may affect signal propagation
- Bruno Bertrand, *Royal Observatory Of Belgium* – next talk


Galileo Satellites

- Hydrogen Masers
- Potentially large improvements from taking advantage of properties of Galileo clocks + satellites
- Francesco Vespe, *Agenzia Spaziale Italiana* – next talk+1

| | | |
|-------|---|---|
| 09:20 | Bertrand <i>Royal Observatory Of Belgium</i> | Fundamental physics tests using the propagation of GNSS signals |
| 09:40 | Vespe <i>Agenzia Spaziale Italiana</i> | Dark Matter Search by GALILEO |

Aside: challenges of re-purposed data



data from JPL: Histogram



- Possible that some clocks mis-identified
(Here, one of the “Rb” clocks is probably Cs).
- Same discrepancy in autocorrelation function, Allan variance etc.

Aside: challenges of re-purposed data

data from JPL: Histogram




- Possible that some clocks mis-identified
(Here, one of the “Rb” clocks is probably Cs).
- Same discrepancy in autocorrelation function, Allan variance etc.

Extra: Asymmetry

Small objects: no correlated signal

- small size $\sim \rightarrow$ large rate
- Shift in mean: unobservable (DM always present)
- Induce non-Gaussian features (such as an asymmetry)




$$\mathcal{R} = \frac{1}{\mathcal{T}} = \frac{3\rho_{\text{DM}} v_g}{4\rho_{\text{inside}} R}, \quad \kappa_3 \approx \frac{2\mathcal{R}\tau_0 \chi_0^3}{5\sigma^3}.$$

Extra: Annual modulation




- Yearly change in event rate:
- Sun + Earth velocities add
- $R(t) = R_0 + R_m \cos(\omega t + \phi_{\text{June}2})$
- $\Delta \kappa_3 / \kappa_3 = 10\%$
- $\Delta \kappa_4 / \kappa_4 = 15\%$



Extra: Limits Q-balls: α (photon field)

Proof of concept: Limits from GPS Cs/H $K_\alpha(Cs/H) \simeq 0.8$



- BMR, Derevianko, arXiv:1803.00617


Red line: sensitivity estimate for 1 year of optical Sr

- Can also place limits on topological defects

Extra: Search method

Fibre network

- Max-likelihood fit method¹
- Coherent α variations
- $N_{\text{pairs}} \geq 2$ - to use network




$$p(D_t | \delta\alpha, \theta) = C p(\theta) \exp \left(\frac{-1}{2} [d - \delta\alpha s]^T H [d - \delta\alpha s] \right),$$

d : data stream; $s = s(\theta)$ signal template, $\delta\alpha$: magnitude; $H = E^{-1}$ covariance

a) Signal-to-noise $R >$ threshold?
⇒ event detection



b) Largest $\delta\alpha$ that cannot be ruled out?
⇒ set limits



$$\delta\alpha^{\text{bf}} = dHs/sHs. \quad \Rightarrow \quad \delta\alpha < dHs/sHs + n(sHs)^{-1/2}.$$

(1) BMR, G. Blewitt, C. Dailey, and A. Derevianko, Phys. Rev. D **97**, 083009 (2018).


Extra: Search results



• arXiv:1907.02661

- With >50 hours of total usable network data
(no gaps, $N \geq 4$) taken over \sim month in 2017:
- No significant events found: place constraints

Extra: Resolve speed + direction



Resolution: simulation using GPS

- Resolve v – DM vel. distro is “known” – reject false positives!
- Many clocks + High sampling frequency and/or **Large distances**